

Socket Shield Technique- A novel approach for esthetic region of the jaw

¹Dr. Shreya Gupta, M.D.S, Assistant Professor, Dept. of Prosthodontics, Trivani Dental College, Bilaspur.

²Dr. Anubhuti Sabhlok, M.D.S, Hitkarini Dental College and Hospital, Jabalpur.

³Dr. Muzammil, M.D.S, Assistant Professor, Dept. of Periodontology, College of Applied Health Sciences, Al Rass, Qassim University, Buraidah.

⁴Dr. Kajal Talele, B.D.S, M.H.A, Rutgers, The State University of New Jersey.

⁵Dr. Srivastav Aatish, M.D.S, Reader, Dept. of Prosthodontics, Shree Sai Dental College & Research Institute, Andhra Pradesh.

⁶Dr. Ritika Kataria, M.D.S, Dept. of Oral & Maxillofacial Surgery, Himachal Institute of Dental Sciences, Paonta Sahib.

Corresponding Author: Dr. Ritika Kataria, M.D.S, Dept. of Oral & Maxillofacial Surgery, Himachal Institute of Dental Sciences, Paonta Sahib.

Citation of this Article: Dr. Shreya Gupta, Dr. Anubhuti Sabhlok, Dr. Muzammil, Dr. Kajal Talele, Dr. Srivastav Aatish, Dr. Ritika Kataria, "Socket Shield Technique- A novel approach for esthetic region of the jaw", IJDSIR- February - 2021, Vol. – 4, Issue - 1, P. No. 107 – 112.

Copyright: © 2021, Dr. Ritika Kataria, et al. This is an open access journal and article distributed under the terms of the creative commons attribution noncommercial License. Which allows others to remix, tweak, and build upon the work non commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Type of Publication: Review Article

Conflicts of Interest: Nil

Abstract

Tooth extraction leads to a cascade of events resulting in resorption of alveolar bone around the socket. The buccal bone loss occurring postextraction requires reconstruction to achieve esthetically pleasing results. In the socket-shield technique, the root is bisected and bucca two-third of the root is preserved in the socket, leaving the periodontium along with bundle bone and buccal bone intact. This article aims to collect and evaluate the available literature associated with the socket shield technique as first described by Hurzeler et al (2010).

Keywords: Socket Shield, Immediate Implant, Bone Graft, Osseointegration

Introduction

Dental implants have become a gold standard treatment modality for tooth replacement. Earlier, implants were used to support multi-unit prosthesis, with advancements these are now commonly used to replace single missing tooth, particularly in the aesthetic region. With ever increasing demand to achieve cosmetically pleasing results, it has become mandatory to preserve buccal hard and soft tissues, preventing negative cosmetic impacts. Various authors have suggested that if a tooth root is left in the alveolar socket, resorption of bundle bone is lesser, preserving the buccal bone.^[1] However, deliberate root retention to preserve bone is not been practiced in a routine practice, even till date.^[1]

Principle of Socket Shield Technique^[2]

The aim of the socket shield technique is to safeguard the buccal plate by creating a buccal shield. Tooth marked for extraction is not extracted completely, the root is sectioned in a fashion that the facial root section remains in its physical relationship to the buccal plate, which in future would prevent alveolar bone loss.

Table 1: Indications and Contraindications^[2]

Indications	Contraindications
For prevention OF Buccal plate in case of immediate implants	Loss of buccal bone due to vertical fracture
In vertically fractured tooth without pupal pathologies.	Loss of buccal bone due to periodontitis.
To secure dental papilla between dental implants	Root caries

Table 2: Advantages And Disadvantages^[3]

Advantages	Disadvantages
No additional material cost	Not yet predictable
No co-morbidity	No long term data available
Single surgery	Technique sensitive
Can be done in sites with apical pathology	Require more time
Assures safeguard of peri-implant tissues.	Shifting of buccal root portion or even buccal lamellar bone.
Helps to retain esthetics	
Guides to place implant in proper location	
Complete osseointegration can be achieved	

Steps of Surgical Procedure^[3]

Under local anesthesia, following procedure is done:

- Clinical crown is separated above the gingival level, with a diamond bur.
- Using long tapered fissure diamond bur, tooth is sectioned vertically.
- The palatal root portion is removed carefully with periostomes and forceps, without putting stress on the buccal tissue.
- Root portion is thinned to a thickness of 2mm using round diamond bur with saline irrigation.
- An osteotomy is planned and implant is placed palatal to the socket shield.
- Gap between the shield and implant is left to permit blood clot formation.
- Implant site is then closed with sutures.

Discussion

Hurzeler et al.^[1,4] in his study (in beagle dogs) published a proof-of-principle study stating that leaving a 1.5 mm thick root fragment on the buccal aspect of the implant site, would be sufficient leaving space for implant placement, maintaining the buccal plate.

Figures 1-7 illustrates the socket shield technique by Hurzeler et al.^[1]

A recently published systematic review by Mourya et al.^[5] stated that modifications to the technique gave promising positive results, although the graft application did not play much role associated with socket shield technique. Thereafter, a study by Christian et al. also does not clearly shows whether the socket shield technique will provide a stable long term outcome.^[1]

Baumer et al.^[6] stated in there study that the socket shield technique shows great esthetic results with preservation of facial tissue contour. Likewise, Siormpas KD et al.

published in their study that intentional root retention with its intact periodontal mechanism can lead to continuous osseointegration of implants placed in maxillary anterior region.^[7]

Kan et al. observed in a case with modified shield technique, with the shield located in the interproximal areas for inter-implant papilla preservation with a successful outcome in preserving bone level and periodontium.^[8]

Glocker et al. performed three cases of modified shield technique and delayed implant placement. After six months the new bone forming and the residual ridge was assessed and preserved via this technique.^[9]

Likewise, similar outcome was reported by Chen et al,^[10] Mitsias et al,^[11] Lagas et al,^[12] Engelke et al,^[13] Al Dary et al.^[14]

Anas et al. reported SS technique to be fruitful in esthetically challenging areas.^[3]



Figure3: Socket Shield invivo




Figure 4: Implant placed



Figure 5: Healed implant site



Figure 6: Implant restoration in situ



Figure 1: Remaining root section




Figure 2: Socket Shield schematic (transverse view)




Figure 7: Postoperative X-Ray after Osseo integration

Classification^[15]




Figure 8: Type 1




Figure 9: Type 2




Figure 10: Type 3




Figure 11: Type 4



Figure 12: Type 5




Figure 13: Type 6

Classification of Predicted Height of Interdental Papilla^[15]

The classification is proposed to help in understanding the clinical application of this technique (Fig.8-13).

Type 1: Buccal shield

The shield lies in the buccal part of the socket (between proximal line angles of tooth).

It is indicated in single edentulous site with both mesial and distal tooth present.

Type 2: Full C Buccal shield

The shield lies in buccal part and the interproximal part on both the sides of the socket.

It is indicated for the following clinical scenarios:

- Implant existing on either side of the proposed site.
- Missing tooth on either side without an implant.
- Having implant on one side and missing tooth on one side.

Type 3: Half C Buccal shield

The shield lies in buccal part and the interproximal part.

It is recommended in case of tooth on one side and implant on the other side.

Type 4: Interproximal shield

Here the shield lies only in the mesial/distal part of the socket.

It is indicated in case of buccal bone resorption, requiring graft with implant/missing tooth on a side.

Type 5: Lingual-palatal shield

The shield lies on the lingual/palatal side of the socket.

It can be considered for maxillary molars.

Type 6: Multiple Buccal shields

It is a case with two or more shields in the socket.

It is indicated in cases with a vertical root fracture.

Complications^[1]

Socket shield exposure as reported by Gluckman et al.^[16] was the most commonly reported with a total of 17 exposed socket shields. They reported 12 internal and 4 external shields exposed. Two of the external exposures required a connective tissue graft for closure and three infected socket shield needed to be removed, however the implants were retained.

Various other complications reported were:

- Resorption of the socket shield
- Peri-implantitis
- Non-integration of the implants

Conclusion

Socket shield technique is gaining popularity among the clinicians worldwide. It has proven to be a promising technique for the preservation of bony buccal plate for immediate implant placement.

References

1. Hurzeler MB, et al. The socket-shield technique: a proof-of-principle report. J Clin Periodontol. 2010;37(9):855–62.
2. Gluckman H, Du Toit J, Salama M. The socket-shield technique to support the buccofacial tissues at immediate implant placement. Int Dentistry. 2015;5:614.
3. Anas B, Shenoy KK. Socket Shield Technique-A Neoteric Approach in Ridge Preservation. Sch. J. Dent. Sci., 2017; 4(3):125.

4. Hurzeler MB et al. The socket-shield technique: a proof-of-principle report. *J Clin Periodontol* 2010; 37: 855–862.
5. Mourya A, et al. Socket-shield technique for implant placement to stabilize the facial gingival and osseous architecture: a systematic review. *J Investig Clin Dent*. 2019;10(4):e12449.
6. Baumer D, Zuhr O, Rebele S, Schneider D, Schupbach P, Hurzeler M. The socket-shield technique: first histological, clinical, and volumetrical observations after separation of the buccal tooth segment - a pilot study. *Clin Implant Dent Relat Res*. 2015;17(1):71-82.
7. Siormpas KD, Mitsias ME, Kontsiotou-Siormpa E, Garber D, Kotsakis GA. Immediate Implant Placement in the Esthetic Zone Utilizing the “RootMembrane” Technique: Clinical Results up to 5 Years Post loading. *International Journal of Oral & Maxillofacial Implants*. 2014 Dec 1;29(6).
8. Kan JY, Rungcharassaeng K. Proximal socket shield for inter implant papilla preservation in the esthetic zone. *International Journal of Periodontics & Restorative Dentistry*. 2013 Jan 1;33(1).
9. Glocker M, Attin T, Schmidlin PR. Ridge preservation with modified “socket-shield” technique: a methodological case series. *Dentistry Journal*. 2014 Jan 23;2(1):11-21.
10. Chen CL, Pan YH. Socket Shield Technique for Ridge Preservation: A Case Re-port. *J Prosthodont Implantol*. 2013;2(2):16-21.
11. Mitsias ME, Siormpas KD, Kontsiotou-Siormpa E, Prasad H, Garber D, Kotsakis GA. A Step-by-Step Description of PDL-Mediated Ridge Preservation for Immediate Implant Rehabilitation in the Esthetic Region. *The International journal of periodontics & restorative dentistry*. 2015;35(6):835-41.
12. Lagas LJ, Peplinkhuizen JJ, Bergé SJ, Meijer GJ. Implanteren in de esthetische zone: de socket shieldtechniek. *Ned Tijdschr Tandheelk*. 2015; 122:33- 6.
13. Engelke W, Beltrán V, Decco O, Valdivia-Gandur I, Navarro P, Fuentes R. Changes in morphology of alveolar buccal walls following a traumatic internal root fragmentation.
14. Al Dary H, Al Hadidi A. The Socket Shield Technique using Bone Trepine: A Case Report. *International Journal of Dentistry and Oral Science*. 2015 Nov 5;5:1- 5.
15. : Kumar PR, Kher U. Shield the socket: Procedure, case report and classification. *J Indian Soc Periodontol* 2018;22:266-72.
16. Cherel F, Etienne D. Papilla preservation between two implants: a modified socket-shield technique to maintain the scalloped anatomy? A case report. *Quintessence Int*. 2014;45(1):23.