

Immunological Aspects of Periodontal Diseases – A Review

¹Raina JP Khanam, PG Student, Department of periodontology, Himachal institute of Dental Sciences, Paonta Sahib, Himachal Pradesh

²Rajan Gupta, Head and professor, Department of periodontology, Himachal institute of Dental Sciences, Paonta Sahib, Himachal Pradesh

³Deepti Shakya, PG Student, Department of periodontology, Himachal institute of Dental Sciences, Paonta Sahib, Himachal Pradesh

⁴Kanika Thakur, PG Student, Department of periodontology, Himachal institute of Dental Sciences, Paonta Sahib, Himachal Pradesh

⁵Shilpa Kaundal, PG Student, Department of periodontology, Himachal institute of Dental Sciences, Paonta Sahib, Himachal Pradesh

⁶Tenzing Yutso Bhutia, PG Student, Department of periodontology, Himachal institute of Dental Sciences, Paonta Sahib, Himachal Pradesh

Corresponding Author: Raina JP Khanam, PG Student, Department of periodontology, Himachal institute of Dental Sciences, Paonta Sahib, Himachal Pradesh

Citation of this Article: Raina JP Khanam, Rajan Gupta, Deepti Shakya, Kanika Thakur, Shilpa Kaundal, Tenzing Yutso Bhutia, “Immunological Aspects of Periodontal Diseases – A Review”, IJDSIR- July - 2021, Vol. – 4, Issue - 4, P. No. 288 – 297.

Copyright: © 2021, Raina JP Khanam, et al. This is an open access journal and article distributed under the terms of the creative commons attribution noncommercial License. Which allows others to remix, tweak, and build upon the work non commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Type of Publication: Review Article

Conflicts of Interest: Nil

Abstract

Periodontal disease is recognized as a major public health problem throughout the world and is the most common cause of tooth loss in adults. Although periodontal disease is of microbial etiology, the determination that periodontal tissue destruction is primarily due to the host response, has created areas of research directed at altering an individual’s reaction to the bacterial challenge. This article will be focusing on various aspects of immunology in periodontal diseases such as immunity itself, cells of the

immune system, leukocyte functions, molecular biology, inflammatory mediators of the immune system, osteoimmunology and immunology in various periodontal diseases.

Keywords: Immunity, periodontal diseases, host-microbial, osteoimmunology, leukocyte.

Introduction

Periodontal disease is a chronic bacterial infection that affects the gingiva and bone that supports the teeth. There are some immunological factors involved in the

development and control of this oral disease, such as: the participation of inflammatory cells in local inflammation, the synthesis of chemotaxis proteins with activation of the complement system and a range of antimicrobial peptides, such as defensins, cathelicidin and saposins.

An individual's susceptibility to periodontitis may be related to whether plasma cells predominate in the tissues of an individual, or a site, in response to the microbial insult from dental plaque. Various features that play role in immunological responses may include homing of immune and inflammatory cells to target tissues; their local proliferation and synthetic activity; the cytokine profile; and the immunoglobulin subclasses of locally produced antibodies.

Role of innate immunity in periodontal diseases

Defenses against infection include a wide range of mechanical, chemical and microbiologic barriers that prevent pathogens from invading the cells and tissues of the body. Saliva, Gingival Crevicular Fluid and the epithelial keratinocytes of the oral mucosa all protect the underlying tissues of the oral cavity and the peridontium. The commensal microbiota (e.g., in dental biofilm) may also be important for providing protection against infection by pathogenic microorganisms through effective competition for resources and ecologic niches and also by stimulating protective immune responses.

Aspects of innate immunity that are relevant to periodontal disease are now considered.

Epithelial Tissues

The epithelial tissues play a key role in host defense because they are the main site of the initial interactions between plaque bacteria. The keratinized epithelium of the Sulcular and gingival epithelial tissues provides protection for the underlying periodontal tissue in addition to acting as a barrier against bacteria and their products.^[19,20]

Saliva

The action of shear forces associated with saliva flow is important for preventing the attachment of bacteria to the dentition and oral mucosal surfaces. Human saliva also contains numerous molecular components that contribute to host defenses against bacterial colonization and periodontal disease such as Antibodies (e.g., immunoglobulin A), Histatins, Cystatins, Lactoferrin, Lysozyme, Mucins and Peroxidase.

Gingival Crevicular Fluid

GCF originates from the post-capillary venules of the gingival plexus. It has a flushing action in the gingival crevice, but it also likely functions to bring the blood components (e.g., neutrophils, antibodies, complement components) of the host defenses into the sulcus.^[19] The flow of GCF increases in inflammation.^[23]

One of the primary challenges of the innate system is the discrimination of pathogens from host. This challenge is met by recognition of the evolutionary structures: The PRRs (Pattern Recognition Receptors) that bind Pathogen-Associated Molecular Patterns (PAMPs), found in a broad type of organisms.

On the basis of function PRRs are classified as:

Signaling PRRs

1. Toll-like receptors
2. NOD receptors (Nucleotide-Oligomerization domain)

Endocytic PRRs

1. CLRs (C-type lectin receptors)
2. RLRs (RIG-1 like receptors)

On the basis of location PRRs are classified as:

1. Membrane bound PRRs
2. Cytoplasmic PRRs
3. Secreted PRRs

Toll-Like Receptors

Toll-like receptors are the gate keepers of innate immunity. TLRs can be divided into five subfamilies:

TLR2, TLR3, TLR4, TLR5 and TLR9. TLRs are expressed in the peridontium in health and disease. Both commensal and pathogenic periodontal bacteria stimulate TLR-2 signaling.^[35,36] Over-production of pro-inflammatory cytokines due to chronic stimulation of toll-like receptors may lead to tissue destruction.^[29,30]

Lipopolysaccharide Binding Protein/ CD14

The concentration of soluble CD14 receptor in saliva and the systemic level of the soluble form of CD14 is significantly increased in patients with periodontal disease and showed severity dependence with increasing levels of periodontal breakdown. Significantly lower levels of the soluble CD14 protein were observed at sites with advanced attachment loss, indicating a protective effect for CD14.

Nucleotide-binding oligomerization domain –like receptors

Clinical investigations have demonstrated that both NOD1 and NOD2 are expressed in human oral epithelium, gingival fibroblast cells and periodontal ligament.^[42,43,44] In an investigation it was found that mice deficient in NOD2 showed comparable levels of alveolar bone resorption, whereas mice deficient in NOD1 demonstrated reduced levels of alveolar bone loss when compared with wild-type control mice.^[45]

Role of Complement System In Periodontitis

In the context of periodontal inflammation, complement subversion appears to play a major role in periodontal pathogenesis.^[48,49] The dysregulation of complement activities may lead to a failure to protect the host against pathogens and amplify inflammatory tissue damage.^[50,51] Activated complement components are found at higher levels in the gingival crevicular fluid of periodontitis patients as compared with healthy subjects.^[52-54] Local complement activation may promote periodontal inflammation predominantly via C5a-induced

vasodilation, increased vascular permeability and flow of inflammatory exudate, and chemotactic recruitment of inflammatory cells, especially neutrophils.^[55,50]

Role of neutrophils in periodontal diseases

Neutrophils are thought to be key players in host-mediated inflammatory tissue injury in periodontitis and can be found in great numbers in the gingival crevice ($\geq 95\%$ of total leukocytes).^[56] Extravasating neutrophils enter the gingival crevice through the junctional epithelium which, under inflamed conditions, is largely occupied (by about 60%) by trafficking neutrophils. Neutrophils are attracted to infected periodontal tissues by chemoattractants released from bacteria, host cells or degraded tissue. The number of neutrophils increases from 7×10^4 to $20 \times 10^4/\text{ml}$ during the conversion of a healthy sulcus into a diseased gingival pocket.

Neutrophil granules are generally classified into azurophilic (primary), specific (secondary) and gelatinase (tertiary) types. The azurophilic granules contain hydrolytic neutral enzymes such as elastase, cathepsin G, urokinase, myeloperoxidase, lysozymes and mannosidase. The specific granules contain lactoferrin, neutrophil collagenase and lysozymes, whereas the main component of the gelatinase granules is the gelatinocytic MMP-9 (matrix metalloproteinase-9). In addition to these types of granules, a fourth type of granule denoting secretory vesicles is present in mature neutrophils.

Neutrophils abnormalities can lead to congenital defects such as (leukocyte adhesion deficiency, Chediak-Higashi syndrome, Papillon-Lefevre syndrome and chronic/cyclic neutropenia) of the host or by immunosuppressive agents used to treat other systemic diseases, environmental and behavioral factors (smoking) or a variety of strategies designed by the pathogen itself to avoid the protective mechanism of the innate system.

Role of mast cells in periodontal diseases

Mast cells are key elements in the innate immune system and are located throughout the body in close proximity to epithelial surfaces, near blood vessels, nerves and glands, placing them at strategic location for detecting invading pathogens. Mast cells express a number of receptors that allow them to recognize diverse stimuli. In sensitized individuals, IgE is bound to Fcε receptors (fragments crystallizable epsilon receptor) expressed on the mast cell surface and binding of antigen to surface bound IgE induces mast cell activation.

It was revealed in human periodontal disease that an increase in the number of mast cells that may be participating either in the destructive events or in the defense mechanism of the periodontal disease via secretion of cytokines, including perpetuation of the Th2 response and cellular migration and healing processes.

Role of dendritic cells in periodontal diseases

Dendritic cells, including Langerhans cells and dermal dendritic cells are found in gingival tissue and mature CD83+ dendritic cells are present in tissues from patients with periodontitis. It was also reported that *P. gingivalis* and *A. actinomycetemcomitans* stimulated dendritic cells promote a rapid IFN-γ (Interferon gamma) response by stimulating NK cells.

Langerhans cell appears to be principle leukocytes involved in the response of the oral mucosal epithelium to infectious, atopic or dysplastic diseases, whether it be within keratinized (e.g. Gingiva) or non- keratinized (e.g. buccal mucosa) tissue.

Microbial Virulence Factors

The sub-gingival biofilm initiates and perpetuates inflammatory responses in the gingival and periodontal tissues and their primary importance in periodontal pathogenesis is that of activating immune-inflammatory responses that, in turn, result in tissue damage.

➤ Bacterial Enzymes and Noxious Products

These include noxious agents such as ammonia (NH₃) and hydrogen sulfide (H₂S), as well as short-chain carboxylic acids such as butyric acid and propionic acid. These substances have profound effects on host cells. *P. gingivalis* produces two classes of cysteine proteases known as gingipains. They modulate the immune system and disrupt immune-inflammatory responses, thus potentially leading to increased tissue breakdown.^[67]

➤ Microbial Invasion

Periodontal pathogens such as *P.gingivalis* and *Aggregatibacter actinomycetemcomitans* have been reported to invade the gingival tissues,^[71-73] including the connective tissues.^[74] *Fusobacterium nucleatum* can invade oral epithelial cells and bacteria that routinely invade host cells may facilitate the entry of non-invasive bacteria by co aggregating with them.^[75]

➤ Fimbriae

The fimbriae of certain bacterial species, particularly *P. gingivalis* stimulate immune responses, such as IL-6 secretion and the major fimbrial structural component of *P. gingivalis*, FimA, has been shown to stimulate nuclear factor (NF)-κB and IL-8 in a gingival epithelial cell line through TLR-2.^[77,78,79]

➤ Bacterial Deoxyribonucleic Acid and Extracellular Deoxyribonucleic Acid

Bacterial deoxyribonucleic acid (DNA) stimulates immune cells through TLR-9. Extracellular DNA (eDNA) is a ubiquitous constituent of all biofilms and of particular interest in biofilms associated with chronic diseases such as periodontitis.^[83]

Host-Derived Inflammatory Mediators

➤ Cytokines

Cytokines play a fundamental role in inflammation and they are the key inflammatory mediators in periodontal disease.^[88] Cytokines bind to specific receptors on target

cells and initiate intracellular signaling cascades that result in phenotypic changes in the cell by altered gene regulation.^[89,90] Cytokines bind to cell surface receptors and trigger a sequence of intracellular events that lead ultimately to the production of protein by the target cell that alters that cell's behavior. They have profound biologic effects that also lead to tissue damage with chronic inflammation. Cytokines mediate connective tissue and alveolar bone destruction through the induction of fibroblasts and osteoclasts to produce proteolytic enzymes (i.e., MMPs) that break down structural components of these connective tissues.^[91] Cytokines play a key role at all stages of the immune response in periodontal diseases.^[88]

➤ **Prostaglandins**

The prostaglandins (PGs) are a group of lipid compounds derived from arachidonic acid, a polyunsaturated fatty acid found in the plasma membrane of most cells. PGs are important mediators of inflammation, particularly prostaglandin E₂ (PGE₂), which results in vasodilation and induces cytokine production by a variety of cell types. PGE₂ results in the induction of MMPs and osteoclastic bone resorption and it has a major role in contributing to the tissue damage that characterizes periodontitis.

➤ **Matrix Metalloproteinases**

MMPs are a family of proteolytic enzymes that degrade extracellular matrix molecules such as collagen, gelatin and elastin. MMPs are divided into collagenase, gelatinase/type IV collagenase, stromelysins, matrilysins, membrane-type metalloproteinases and others. MMPs are secreted in a latent form (inactive) and are activated by the proteolytic cleavage of a portion of the latent enzyme. MMPs are inhibited by proteinase inhibitors, which have anti-inflammatory properties. Key inhibitors of MMPs found in the serum include the glycoprotein α 1-antitrypsin and α 2-macroglobulin, a large plasma protein produced by

the liver. Inhibitors of MMPs that are found in the tissues include the tissue inhibitors of metalloproteinases (TIMPs), which are produced by many cell types; the most important in periodontal disease is TIMP-1.^[94]

Role of adaptive immunity in periodontal diseases

In addition to the innate immunity, adaptive immunity cells and characteristic cytokines have been described as important players in the periodontal disease pathogenesis scenario. Parts of the adaptive host response in periodontitis as outlined are:

- (1) The nature of the lymphocyte type (T and B cells)
- (2) Antigen recognition by TCRs
- (3) Cytokine profiles of T helper (Th) cells
- (4) Autoimmune reactions that may influence the adaptive host response in periodontitis.

T-Cells In Periodontist

It's evident that both T cells and B cells are present in periodontal disease. It has been hypothesized that T cells are present in more stable lesion of periodontitis while B cells and plasma cells are present in more progressive lesion.

CD4+ T-cells were subdivided initially into two subsets, designated T-helper 1 and T-helper 2, on the basis of their pattern of cytokine production.^[97] T-helper 1 cytokines have been associated with infectious inflammatory bone destruction. T-helper 2 cytokines are described to minimize bone loss.^[98-101] The discovery of new T-helper subsets with prominent roles in the modulation of host responses determined the re-examination of T-helper 1/T-helper 2 dichotomy paradigm in chronic inflammatory diseases, including periodontal diseases.^[102,99,103,104,105]

B-Cells In Periodontist

B cells serve as a well controlled part of the adaptive host response and act on systems regulated by T cells. Different subsets of B cells, such as B-1a and B-2 cells are present in periodontitis lesions. Elevated levels of B-1a

cells have been demonstrated in both periodontitis lesions and peripheral blood of subjects with severe forms of periodontitis. Different pathologic functions of B-1a cells have been associated with IL-10 and this cytokine serves as an autocrine growth factor for this type of B cells.

Osteo-immunology in periodontal diseases

When the response becomes chronic, adaptive immune cells invade the tissue and the inflammatory reaction becomes firmly established, flooding the periodontium with additional bioactive proinflammatory molecular signals (cytokines, chemokines, enzymes, ROS (reactive oxygen species), bacterial products and metabolites, etc.). The accumulation of these molecular signals in the tissue facilitates the spreading of the inflammation to the underlying bone and tamper with the bone homeostasis signaling system, tilting the balance of bone metabolism favoring resorption over formation.

The dynamic behavior of bone enhances its adaptive capacity to functional demands and increases its healing potential after an injury. The balance between bone resorption and apposition is governed by a unified molecular signaling system and is dependent on the effector functions of specialized bone cells. The system that controls the bone metabolism balance comprises the RANKL/OPG/RANK triad, secreted and recognized by the specialized bone effector cells osteoblasts and osteoclasts.

RANKL (receptor activator of nuclear factor κ B ligand) is a cytokine, member of the TNF family that can be membrane bounded or secreted and stimulates osteoclasts differentiation, cell fusion and activation leading to bone resorption through calcium-dependent activation of the transcription of NFATc1 (nuclear factor of activated T-cells 1) gene. RANKL interacts with its cognate receptor RANK in the surface of osteoclast and osteoclasts

precursors, triggering their recruitment to the bone surface, cell fusion and activation.

Osteoprotegerin (OPG) is a soluble protein upregulated in inflammatory conditions, with the capacity to block RANKL's biological functions by competitive inhibition, acting as a decoy receptor, limiting the availability of RANKL able to bind to RANK. A high ratio of RANKL/OPG creates the conditions favorable to bone resorption, while a low RANKL/OPG ratio favors bone apposition. During bacteria-induced inflammation in experimental periodontitis, it has been demonstrated a net increase in the RANKL/OPG ratio, leading to osteoclast genesis and bone resorption. The key event in the inflammatory alveolar bone resorption in periodontitis is the manifold increase in the tissue levels of RANKL unaccompanied by an equivalent increase in OPG levels.

Immunology in Periodontal Diseases

Gingivitis: It is a primary response to the bacteria in plaque. It includes a vascular response with increased fluid accumulation and inflammatory cell infiltration. The early response is mostly lymphocytic, represented by T cells, which is slightly higher. Acute phase protein including α_2 macroglobulin, α_1 anti trypsin and transferrin are increased with gingival inflammation reflecting the locally stressed environment. Result of a study by Kinane et al indicate that IL-1 levels in GCF increase with plaque and peak levels of this mediator precede clinical signs of inflammation in experimental gingivitis.

Chronic Periodontitis

Extensive studies have been performed examining the characteristics of the inflammatory infiltrate in chronic periodontitis. The distribution of gingival mononuclear cells has shown plasma cells (5% to 15%) in chronic periodontitis. The plasma cells are predominated by IgG followed by IgA in tissues from chronic periodontitis. Moreover, IgG cells in the gingival tissues were identified

as IgG1 > IgG2 > IgG3 > IgG4 and IgA1, with high level IgA2 cell levels in advanced lesions. A large number of B cell lineage cells in gingival tissues present a phenotype that can strongly stimulate auto reactive T cells. The T_H/T_S (T helper and T suppressor) ratios are reduced in chronic periodontitis periodontal lesions. The lower CD4/CD8 ratios, low responses to mitogens and higher expression of HLA-DR (human leukocyte antigen DR isotype) on $CD8^+$ T cells have been suggested to be regulators of periodontal progression. Recently, NK cells have been identified in the gingiva of periodontitis (3% to 7%) in association with damaged fibroblasts. Finally, evidence has suggested a depressed functional activity of cells from gingiva in periodontitis in response to mitogenic stimulation. These findings are consistent with an altered immune cell distribution, immunoregulation and/or function that may contribute to the progression of periodontitis.

Extensive studies have been reported that PgE_2 levels are increased in periodontitis when compared to healthy sites with respect to IL-1, investigations have shown:

- 1) IL-1 α and IL-1 β activity in > 70% of GCF.
- 2) IL-1 β levels appear higher in GCF from chronic periodontitis when compared to healthy sites and active sites verses inactive sites, which decrease after treatment.
- 3) IL-1 β and cells producing this cytokine are elevated in tissues of chronic periodontitis and are detected in the lamina propria of these tissues.
- 4) Most studies were unable to document a relationship between IL-1 β and the clinical parameters (ie, probing depth, gingival index, bone resorption) of the site.

Aggressive Periodontist

The prevalence of a humoral immune response to *A.actinomycetemcomitans* is elevated in patients with localized aggressive periodontitis. Numerous mechanisms of serum mediated bacterial killing are proposed,

including lysis by the membrane attack complex of complement and antibacterial substances such as lysozymes. Generalized aggressive periodontitis is often characterized by defects in either neutrophils or monocytes.

There are two neutrophilic function defects i.e. impaired neutrophil function and hyperactive neutrophil function. Selectins and Integrins play a key role in the initial adhesion of neutrophils, thus facilitating transendothelial migration. Peripheral and GCF neutrophil CD18/CD11a and CD18/CD11b expressions have been compared in aggressive periodontitis patients and healthy patients in various studies. It has been demonstrated that neutrophils from patients with aggressive periodontitis have increased intracellular levels of β -glucuronidase, which is present in azurophilic granules of the neutrophils.

Severe periodontal manifestations are also associated with congenital neutropenia, cyclic neutropenia, leukocyte adhesion deficiency type I and type II, glycogen storage disease, Ehlers–Danlos syndrome and Cohen syndrome.^[147] There is also a report of aggressive periodontitis associated with Fanconi anemia.^[148] Fanconi anemia is an autosomal recessive disorder affecting all bone marrow elements and is associated with cardiac, renal and limb malformations as well as with dermal pigmentary changes.

Summary and conclusion

Present knowledge favors the concept that oral bacteria and their metabolic products are the major etiologic factors in the pathogenesis of periodontal disease, although the exact nature of this host-parasite interaction is still incompletely understood. However, a great deal of evidence now suggests that the host's immunologic responses to microbial products are primarily responsible for the development of this disease.

In periodontal disease, the inflammatory response causes tissue resistance to bacterial invasion but also provides mechanisms that contribute to tissue damage. The acute inflammatory cells, such as mast cells, macrophages, dermal dendrocytes, Langerhans cells and polymorphonuclear leukocytes (PMNs), all combine to form a potent anti-bacterial defense mechanism, but the initial signs of gingival inflammation represent the tissue destructive aspects of these cells. In established periodontal disease, there is also chronic inflammatory change, with B cells and T cells adding to the anti-bacterial spectrum. These lymphocytic cells also have the capacity to release cytokines, which may induce the synthesis of arachidonate metabolites (especially PgE₂) and to stimulate macrophages and osteoclasts to release hydrolases and collagenases which are responsible for loss of collagen and bone. RANKL expression is increased in periodontitis tissue compared with healthy gingival tissue. RANKL is expressed on osteoblasts and activated T cells. Both osteoblasts and periodontal ligament fibroblasts stimulated with lipopolysaccharide and IL-1 express RANKL, suggesting that once the inflammation extends into the periodontal ligaments and alveolar bone, alveolar bone resorption might be accelerated through the increased expression of RANKL.

References

1. Bartold PM, Walsh LJ and Narayanan AS. Molecular and cell biology of the gingiva. *Periodontol* 2000; 24:28–55.
2. Schroeder HE and Listgarten MA. The gingival tissues: the architecture of periodontal protection. *Periodontol* 1997; 13:91–120.
3. Ben-Sasson SZ, Hu-Li J, Quiel J, et al. IL-1 acts directly on CD4 T cells to enhance their antigen-driven expansion and differentiation. *Proc Natl Acad Sci USA* 2009; 106:7119–7124.
4. Kikkert R, Laine ML, Aarden LA, et al. Activation of toll-like receptors 2 and 4 by gram-negative periodontal bacteria. *Oral Microbiol Immunol* 2007; 22:145–151.
5. Yoshioka H, Yoshimura A, Kaneko T, et al. Analysis of the activity to induce toll-like receptor TLR2 and TLR4-mediated stimulation of supra-gingival plaque. *J Periodontol* 2008; 79:920–928.
6. Beklen A, Hukkanen M, Richardson R and Konttinen YT. Immunohistochemical localization of Toll-like receptors 1-10 in periodontitis. *Oral Microbiol Immunol* 2008; 23:425-431.
7. Watanabe K, Iizuka T, Adeleke A, Pham L, Shlimon AE, Yasin M, et al. Involvement of toll-like receptor 4 in alveolar bone loss and glucosehomeostasis in experimental periodontitis. *J Periodontal Res* 2011; 46:21-30.
8. Sugawara Y, Uehara A, Fujimoto Y, et al. Toll-like receptors, NOD1 and NOD2 in oral epithelial cells. *J Dent Res* 2006; 85:524–529.
9. Liu J, Wang Y and Ouyang X. Beyond toll-like receptors: *Porphyromonas gingivalis* induces IL-6, IL-8, and VCAM-1 expression through NOD-mediated NF-kappaB and ERK signaling pathways in periodontal fibroblasts. *Inflammation* 2014; 37:522–533.
10. Uehara A and Takada H. Functional TLRs and NODs in human gingival fibroblasts. *J Dent Res* 2007; 86:249–254.
11. Jiao Y, Darzi Y, Tawaratsumida K, et al. Induction of bone loss by pathobiont-mediated Nod1 signaling in the oral cavity. *Cell Host Microbe* 2013; 13:595–601.
12. Hajishengallis G, Liang S, Payne MA, et al. Low-abundance biofilm species orchestrates inflammatory periodontal disease through the commensal microbiota

- and complement. *Cell Host Microbe* 2011; 10:497–506.
13. Maekawa T, Krauss JL, Abe T, et al. *Porphyromonas gingivalis* manipulates complement and TLR signaling to uncouple bacterial clearance from inflammation and promote dysbiosis. *Cell Host Microbe* 2014; 15:768–778.
14. Hajishengallis G, Maekawa T, Abe T, et al. Complement involvement in periodontitis: molecular mechanisms and rational therapeutic approaches. *Adv Exp Med Biol* 2015; 865:57–74.
15. Ricklin D, Hajishengallis G, Yang K, et al. Complement: a key system for immune surveillance and homeostasis. *Nat Immunol* 2010; 11:785–797.
16. Attstrom R, Laurel AB, Lahsson U, et al. Complement factors in gingival crevice material from healthy and inflamed gingiva in humans. *J Periodontol Res* 1975; 10:19–27.
17. Patters MR, Niekrash CE and Lang NP. Assessment of complement cleavage in gingival fluid during experimental gingivitis in man. *J Clin Periodontol* 1989; 16:33–37.
18. Ryu OH, Choi SJ, Linares AM, et al. Gingival epithelial cell expression of macrophage inflammatory protein-1alpha induced by interleukin-1beta and lipopolysaccharide. *J Periodontol* 2007; 78:1627–1634.
19. Hajishengallis G. Complement and periodontitis. *Biochem Pharmacol* 2010; 80:1992–2001.
20. Delima AJ and Van Dyke TE. Origin and function of the cellular components in gingival crevice fluid. *Periodontol* 2000 2003; 31:55-76.
21. Potempa J and Pike RN. Corruption of innate immunity by bacterial proteases. *J Innate Immun* 2009; 1:70–87.
22. Christersson LA, Albini B, Zambon JJ, et al. Tissue localization of *Actinobacillus actinomycetemcomitans* in human periodontitis. I. Light, immunofluorescence and electron microscopic studies. *J Periodontol* 1987; 58:529–539.
23. Hillmann G, Dogan S and Geurtsen W. Histopathological investigation of gingival tissue from patients with rapidly progressive periodontitis. *J Periodontol* 1998; 69:195–208.
24. Saglie FR, Marfany A and Camargo P. Intragingival occurrence of *Actinobacillus actinomycetemcomitans* and *Bacteroides gingivalis* in active destructive periodontal lesions. *J Periodontol* 1988; 59:259–265.
25. Saglie FR, Newman MG, Carranza FA Jr, et al. Bacterial invasion of gingiva in advanced periodontitis in humans. *J Periodontol* 1982; 53:217–222.
26. Edwards AM, Grossman TJ and Rudney JD. *Fusobacterium nucleatum* transports non invasive *Streptococcus cristatus* into human epithelial cells. *Infect Immun* 2006; 74:654–662.
27. Lamont RJ and Jenkinson HF. Life below the gum line: pathogenic mechanisms of *Porphyromonas gingivalis*. *Microbiol Mol Biol Rev* 1998; 62:1244–1263.
28. Ogawa T and Uchida H. A peptide, ALTTE, within the fimbrial subunit protein from *Porphyromonas gingivalis*, induces production of interleukin 6, gene expression and protein phosphorylation in human peripheral blood mononuclear cells. *FEMS Immunol Med Microbiol* 1995; 11:197–205.
29. Asai Y, Ohyama Y, Gen K, et al. Bacterial fimbriae and their peptides activate human gingival epithelial cells through Toll-like receptor 2. *Infect Immun* 2001; 69:7387–7395.

30. Jakubovics NS and Burgess JG. Extracellular DNA in oral microbial biofilms. *Microbes Infect* 2015; 17:531–537.
31. Preshaw PM and Taylor JJ. How has research into cytokine interactions and their role in driving immune responses impacted our understanding of periodontitis? *J Clin Periodontol* 2011; 38:60–84.
32. Birkedal-Hansen H. Role of cytokines and inflammatory mediators in tissue destruction. *J Periodontol Res* 1993; 28:500–510.
33. Taylor JJ, Preshaw PM and Donaldson PT. Cytokine gene polymorphism and immunoregulation in periodontal disease. *Periodontol 2000* 2004; 35:158–182.
34. Bartold PM and Narayanan AS. Molecular and cell biology of healthy and diseased periodontal tissues. *Periodontol 2000* 2006; 40:29–49.
35. Birkedal-Hansen H. Role of matrix metalloproteinases in human periodontal diseases. *J Periodontol* 1993; 64:474–484.
36. Murphy KM and Reiner SL. The lineage decisions of helper T cells. *Nat Rev Immunol* 2002; 2:933-944.
37. Colic M, Gazivoda D, Vucevic D, Vasilijic S, Rudolf R and Lukic A. Proinflammatory and immunoregulatory mechanisms in periapical lesions. *Mol Immunol* 2009; 47:101-13.
38. Graves DT, Oates T and Garlet GP. Review of osteoimmunology and the host response in endodontic and periodontal lesions. *J Oral Microbiol* 2011; 17:3.
39. Queiroz-Junior CM, Silva MJ, Corrêa JD, Madeira MF, Garlet TP, Garlet GP, et al. A controversial role for IL-12 in immune response and bone resorption at apical periodontal sites. *Clin Dev Immunol* 2010.
40. Repeke CE, Ferreira SB Jr, Vieira AE, Silveira EM, Avila- Campos MJ, Silva JS, et al. Dose-response met-RANTES treatment of experimental periodontitis: a narrow edge between the disease severity attenuation and infection control. *PLoS One* 2011; 6.
41. Botero J, Contreras A, Lafaurie G, Jaramillo A, Betancourt M and Arce RM. Occurrence of periodontopathic and superinfecting bacteria in chronic and aggressive periodontitis subjects in a Colombian population. *J Periodontol* 2007; 78:696-704.
42. Jager A and Kuchroo VK. Effector and regulatory T-cell subsets in autoimmunity and tissue inflammation. *Scand J Immunol* 2010; 72:173-184.
43. Silva TA, Garlet GP, Fukada SY, Silva JS and Cunha FQ. Chemokines in oral inflammatory diseases: apical periodontitis and periodontal disease. *J Dent Res* 2007; 86:306-319.
44. Wan YY. Multi-tasking of helper T cells. *Immunology* 2010; 130:166-171.
45. Tang L, Zhou XD, Wang Q, et al. Expression of TRAF6 and pro-inflammatory cytokines through activation of TLR2, TLR4, NOD1, and NOD2 in human periodontal ligament fibroblasts. *Arch Oral Biol* 2011; 56:1064–1072.