

**Shade Selection Blending of Conventional and Digital Methods - An Updated Review**

<sup>1</sup>Bhagyashri Katade, Postgraduate Student, Department of Prosthodontics, Crown & Bridge and Implantology, C.S.M.S.S. Dental College & Hospital, Aurangabad, Maharashtra.

<sup>2</sup>Aishwarya Mundlik, Postgraduate Student, Department of Prosthodontics, Crown & Bridge and Implantology, C.S.M.S.S. Dental College & Hospital, Aurangabad, Maharashtra.

<sup>3</sup>Ruchi Kasat, Professor and Guide, Department of Prosthodontics, Crown & Bridge and Implantology, C.S.M.S.S. Dental College & Hospital, Aurangabad, Maharashtra.

<sup>4</sup>Nazish Baig, Professor and Guide, Department of Prosthodontics, Crown & Bridge and Implantology, C.S.M.S.S. Dental College & Hospital, Aurangabad, Maharashtra.

**Corresponding Author:** Aishwarya Mundlik, Postgraduate Student, Department of Prosthodontics, Crown & Bridge and Implantology, C.S.M.S.S. Dental College & Hospital, Aurangabad, Maharashtra.

**Citation of this Article:** Bhagyashri Katade, Aishwarya Mundlik, Ruchi Kasat, Nazish Baig, “Shade Selection Blending of Conventional and Digital Methods - An Updated Review”, IJDSIR- April - 2022, Vol. – 5, Issue - 2, P. No. 149 – 155.

**Copyright:** © 2022, Aishwarya Mundlik, et al. This is an open access journal and article distributed under the terms of the creative commons attribution noncommercial License. Which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**Type of Publication:** Review Article

**Conflicts of Interest:** Nil

**Abstract**

Major challenges in cosmetic dentistry are to accomplish appropriate and satisfactory reproduction of natural shade of teeth. The procedure of choosing a shade can be performed using visual method or by an instrument. The kind of shade guide, individual ability to choose shades and conditions the choice is made under, all have influence on reliability and accuracy of the procedure. Therefore, it is necessary to introduce instrumental color determination in everyday work .They include RGB devices, colorimeters, spectrophotometers. The impact of the color science can be seen on various restorative materials ranging from ceramics to maxillofacial prosthetic materials. Good communication between the dental team of all these details is paramount to a

successful aesthetic result. It also discusses the colorimetry, which is the science and technology used to quantify and describe physically the human color perception. Theories of color space and clinically portable colorimetric devices developed for determining tooth shade in the dental practice is discussed.

**Keywords:** Tooth color, Shade guide, Spectro photo meter, Colorimeter.

**Introduction**

Dentistry represents a significant portion of dental services encompassing on the blend of science and art. New improvements in technologies in dental materials and consequent aesthetic enhancement of direct and indirect restorations require trained practitioner enabling to choose the right shade. The path to the appropriate

choice of shade for future restoration is not simple. Achieving the satisfying morphological, optical and biological form of a restoration is one of the most important goals of esthetic dentistry and dentistry (1).

Color is defined as subjective perception of the quality of light and colorimetry is a scientific discipline which enables measuring and specifying the color (2). Basic knowledge about shades is prerequisite to make the right choice. In 1900, American painter and teacher of art Albert Henry Munsell wrote about tri-dimensional property of color and parameters such as: hue, chrome and value or lightness (3).

Shade selection procedure in cosmetic dentistry can be done by visual or instrumental color determination. Visual color determination considers comparing the shade with already known physical standard accepted as a reference. Basically, it is the use of color atlas or shade guide in more or less controlled conditions (4). The most common and widely accepted way to choose shade is by the shade guide. The first shade guide “Tooth Color Indicator”, made by Clark, and contained 60 porcelain samples of shades. Years back “Vitapan Classic Shade Guide” was the gold standard in choosing shades. Although still in use today, it shows various shortcomings successfully overcome in new guides (5,6). Hall (1991) work was base for the development of “Vitapan 3D-Master Shade Guide” which was the first commercial system for determining and reproduction the shades based on the principles of colorimeter classification.

Vitapan 3D-Master Shade Guide is a highly improved compared to the classic shade guides with better organization, wider range and uniform distribution of shades (5,6). A new version of VITA Toothguide 3D-Master called “Linearguide 3D Master” is also available. It has a practical linear structure of shades more suitable

for dentists who use Vitapan Classic shade guide which also has a linear structure of shades (5). The instrumental color determination requires devices such as spectro photometers, tri-stimuli colorimeters, spectroradiometers as well as digital cameras (7). Tam et al demonstrated that digital cameras could be substitutes for contact type instruments in shade selection. They proposed a new method to compare the color of shade tabs taken by a digital camera using appropriate color features. Recently, Kim et al developed a polarization dental imaging modality (PDIM) to obtain cross-polarized images and, therefore, to address the problem of specular reflection.

Shade selection procedure depends on various factors including translucency, contour and surface texture. Tooth shade selection using a conventional means involves a high degree of subjectivity. Traditional shade guides are available that use several methods for quantifying shade. Technology-based systems provide with an advantage of natural looking restorations. They include RGB devices, colorimeters, and spectro photo meters.

This article discusses the different selection procedures involved for correct shade matching, the various systems available for the restorative dentist, and the recent developments in the field.

### **Shade matching in dentistry**

Teeth are often termed “polychromatic” and have the variation in hue, value, and chroma within the teeth and give three dimensional depth and characteristics. Christopher (8) stated that the difficulty of shade selection is that clinicians must be able to interpret a multi-layered structure of varying thickness, opacities, and optical surface characteristics. This can affect the way that the eye perceives color. He stated that a number of related factors in selecting shades must also be

understood to achieve a successful result. These factors includes (Winter 1990) translucency, contour, surface texture, luster, and fluorescence. The various Color perception elements are, the light source, the tooth, including textures and layers, the environment, the receiver (eye). Llena et al investigate the intra and inter-device reliability of two tooth color measurement devices: Easy Shade (ES) and Spectro Shade Micro (SSM) and showed that both Easy Shade (ES) and Spectro Shade Micro (SSM) show excellent repeatability, so they can be used in office to evaluate tooth color or to assess color changes after treatment.

### **Measurement of color**

Color determination in dentistry can be divided into two categories, visual and instrumental.

### **Visual technique**

The most frequently used technique in the shade matching of teeth to restorative materials is done manually and visually with dental shade guides. Using this method, shade matching is accomplished by visually comparing a tooth with multiple standards, usually represented as shade guide tabs, Figure 1. The shade selection done in this fashion is subjective and is influenced by the ability of the human observer to choose the best matching shade. Factors such as observer color perception, observer bias, ambient lighting, and the acceptance threshold of mismatch have effects on accurate shade matching. Also, not all commercial shade guides cover all the ranges of value, hue, and chroma present in human tooth structure. [9] The aging of these shade guides combined with potential “darkening” of the tabs due to cold sterilization create havoc if undiscovered. The original shade guides represented only about 30 percent of the color range of natural dentition. This range has improved tremendously with the advances in colorimetry and improved shade

guides. Quite often this creates the need for “cross-matching” which requires the laboratory technologist to reformulate the material shades to match the particular dentition. This is an extremely subjective process which could lead to unacceptable mismatches. Another problem is the influence of shade designation on the difference in color in various brands of dental restorative materials.

The most popular shade guides include the vitapan classical shade guide, Vita 3D master shade guide system, and the chromascop shade guide system. The shade tabs arrangement in the Vita Classical is by hue whereas in the Chromascop guides (10), the tabs are arranged in five clearly discernible value levels. VITAPAN introduced in 1956, the current American Dental Association gold standard for monitoring of tooth whitening, is supposed to correspond to decreasing lightness (from left to right). A very popular shade guide where in tabs of similar hue are clustered into letter groups A (redyellow), B (yellow), C (grey), D (redyellow-gray), and chroma designated with the numerical values (e.g. A1). The manufacture protocols include hue selection followed by chroma and value effect.

Classical shade guide introduced by Miller (11) was too low in chroma and too high in value when compared to natural extracted tooth samples. The “value scale” is inaccurate in terms of decreasing L\* values, exhibiting redundancy and uneven lightness differences among neighboring tabs, which to a certain extent compromises the results of clinical studies that have been performed so far (9).

Bayindir et al (12) stated that the Vitapan 3D master shade guide system results in lower coverage errors than the Vita lumin or Chromascop shade guide systems. Ahn et al. (13) concluded that the color distribution of the

Vitapan 3D master shade guide was more ordered than previously reported color distributions of other, traditional shade guides. However, the interval in the color parameters between adjacent tabs was not uniform. Christopher et al (8) suggested the shade selection sequence to be as follows

- Make sure teeth are clean and unstained before attempting shade selection.
- Shade selection should be completed before preparation as teeth can become dehydrated and result in higher values.
- Shades should be done when the dental team is not fatigued as in the end of the day.
- Ensure surgery surroundings are of neutral color so that there is no color cast onto the teeth.
- Remove lipstick; ask patients not to wear lurid clothing or any items that may distract the attention of the teeth.
- Patient should be in an upright position at a level similar to the operator and the shade guide should be at arm's length. This ensures that the most color sensitive part of the retina will be used.
- Observations should be made quickly (5 seconds) to avoid fatiguing the cones of the eyes. If longer than this, the eye cannot discriminate and the cones become sensitized to complement the observed color.
- Blue fatigue can accentuate yellow sensitivity so dentists can look at a blue object, bib, etc, while resting the eyes.
- Use color corrected light illumination, which should be of a diffuse nature.
- Choose basic shade at the middle of the tooth - using the Vita System 3D-Master technique of value, chroma then hue. Use blue card to avoid chromatic adaptation.

- Viewing tabs through half-closed eyes can decrease ability to discriminate color but increases the ability to match value. Look at the other parts of the teeth, dividing the teeth into 9 sections from apical to incisal, and mesial to distal.
- Necks of shade tabs often can be removed as they have a great deal of colorants that may introduce errors.
- Examine tooth for translucency and any characterizations, e.g. craze line, hypocalcification, etc.
- Create a shade/chromatic map – divided into different sections to ensure correct placement of different effects, characterizations and shades.
- Photograph teeth and tabs using different lighting conditions to minimize metamerism, e.g. flash (5500K) and natural daylight (6500K).
- Photograph teeth at a 1:1 ratio for detailed characterizations.
- Send digitized images and shade map to ceramist.

Conventional visual tools used to determine shade are highly susceptible to various optical illusions and contrast effects. Due to the errors with the use of commercial shade guides, many different devices and machine tools are used in order to make the color assessment more simple, rapid, precise, and perfect. The semi-translucent structure, small size, and irregular surface of teeth contribute to the complexity of this procedure. Several clinical studies have confirmed that computer assisted shade analysis is more accurate and more consistent compared with human shade assessment. The advantages are no influence of surroundings or lighting and the results being reproducible (14).

Different types of technological shade systems are: RGB devices, digital cameras, spectro photo meters, colorimeters.

### **RGB devices**

RED, GREEN, BLUE image information to create a color image. They do not control key variables associated with accurate color determination (14). ShadeScan™ measures shades over the entire tooth surface, then analyzes them and generate a shade match report. It likewise can generate a report to be used with any standard shade guide system. ShadeScan™ creates an image of the tooth with a translucency and characterization map, and then will generate a printed report.

### **Digital cameras**

Digital cameras are efficient and easy to use and can be an ideal supplement for the clinician and lab technician in quantifying shade but alone not a very reliable method for shade analysis (14). Factors such as illumination and the angle of the photograph will alter how color is perceived by the camera. Alvin et al. (15) stated the use of Commercial SLR cameras when combined with the appropriate calibration protocols showed potential for use in the color replication process. Spear stated the use of color-corrected professional quality film (e.g. Kodak EPN-100, E100-S, or EPP) and has a good photo lab to develop them, taking vector shots at 65-70° looking down with an incisal edge away from chroma and hue helps in increasing the amount of reflection (16).

### **Spectrophotometers**

It measures and records the amount of visible radiant energy reflected or transmitted by an object one wavelength at a time for each value, chroma, and hue present in the entire visible spectrum (17, 18). VITA Easyshade Compact is the device that meets the greatest number of requirements for choosing the shades in clinical settings. The device can be used to determine an overall tooth shade, the shade of each third of the tooth-

cervical, middle and incisal, as well as to confirm the shade of the restoration.

### **Colorimeters**

Colorimeters also process reflected light from the surface of an object, and these devices calculate the light flux data into color parameters that can be used to determine shade and calculate color differences.(19) “Color space” is a specific term, identifying a particular combination of color model and mapping function, it tends to be used informally to identify a color model. They provide measurements in CIELAB units (L\*, A\*, B\*) that can compare the color parameters of different objects when analyzed mathematically. Colorimeters can be of two types mainly the photoelectric tristimulus colorimeters (Microcolor) and silicon photodiode array (Orient Scientific Ltd). Microcolor colorimeter (a photoelectric tri-stimulus colorimeter) is a compact color measuring instrument that is less prone to overheating and is cost effective (20). The shade vision system adds the advantage of shade information being sent to the dental laboratory via e-mail, disk, or by printout. The ShadeVision unit uses advanced colorimetry to scientifically determine the hue, value and chroma of the teeth.

### **Stump shade selection**

It is important to communicate the prepared tooth or “stump” shade to the ceramist so that they can build the restoration with the right opacity/ translucency (21). It may be necessary to use a more opaque ceramic to block out discoloration, e.g. an alumina- or zirconia based restoration may be a better choice than a glass-based ceramic.

### **Comparison of Manual and Mechanical Shade Selection Techniques**

As previously mentioned, the manual, visual shade guide systems continue to be the most common means of

shade selection in dentistry. The visual systems are much cheaper to obtain. However, shade selection is very subjective and is heavily influenced by ambient conditions. With the addition of today's advanced colorimetry technology, qualitative and quantitative reliability more than doubles to approximately 70 percent of the true color range of natural dentition. Recently better results were reported with a dental spectrophotometer than the visual method [22,23]. One study [24] compared the repeatability and inter-device reliability between a colorimeter (ShadeEye NCC®) and a spectrophotometer (VITA EasyShade Compact®). Another study which compared the reliability and accuracy of four clinical shade matching instruments (two clinical spectrophotometers and two colorimeters) [25] found variations in both reliability and accuracy of the devices. Most device comparisons demonstrated high reliability which would indicate predictable shade values when repeating the measurements. It has been recommended that both visual and mechanical shade selection system be incorporated into the contemporary dental practice as they complement each other.

### Conclusions

Understanding the influence of different variables in shade selection from light illumination to the tooth's hue, value and chroma and how the eye interprets this can assist in this selection. The use of the Vita System 3D-Master allows a logical selection of color into hue, value and chroma. There are limitations of shade guides as they fail to account for the variability found in natural teeth, e.g. fluorescence, opalescence, translucency, enamel thickness, and objectivity. Effects of surface texture on light reflection and different characterizations must be recorded and duplicated in the final restorations. The use of technology with different devices in shade selection may eliminate subjectivity of choosing and the

use of photography to communicate shades and characterizations has improved the selection process. A procedure of shade selection has been described to ensure consistent results considering the different variables that influence shade matching.

### References

1. Bayindir F, Kuo S, Johnston W, Wee A. Coverage error of three conceptually different shade guide systems to vital unrestored dentition. *J Prosthet Dent* 2007;98:175-85.
2. Milićević V. Sistemi boja u stomatologiji. *Stom Protet* 1998;3:101-05.
3. Curd FM, Jasinevicius RT, Graves A, Cox V, Sadan A. Comparison of the shade matching ability of dental students using two light sources. *J Prosthet Dent* 2006;96:391-96.
4. Hahhad HJ, Jakstat HA, Arnetzl G, Borbely J, VichiA, Dumfahrt H, et al. Does gender and experience influence shade matching quality? *J Dent* 2009;37:40-44.
5. Paravina R. Performance assessment of dental shade guides. *J Dent* 2009;37:15- 20.
6. Marucci B. A shade selection technique. *J Prosthet Dent* 2003;89:518-21.
7. Pusateri SK, Brewer JD, Davis EL, Wee AG. Reliability and accuracy of four dental shade matching devices. *J Prosthet Dent* 2009;101:193-199.
8. Paravina RD, Powers JM. *Esthetic Color Training in Dentistry*. St. Louis MO, USA: Elsevier Mosby; 2004.
9. Brewer JD, Wee A, Seghi R. Advances in color matching. *Dent Clin North Am* 2004;48:341-358
10. Miller LL. *A Scientific approach to shade matching*. Chicago: Quintessence;1980.
11. Bayindir F, Kuo S, Johnston WM, Wee AG. Coverage error of three conceptually different shade



guide systems to vital enrestored dentition. *J Prosthet Dent* 2007;98:175-85.

12. Ahn JS, Lee YK. Color distribution of a shade guide in the value, chroma, and hue scale. *J Prosthet Dent* 2010;100:18- 28.

13. Chu SJ, Devigus A. *Fundamentals of Colors*. Chicago: Quintessence publishing. 2004

14. Wee AG, Lindsey DT, Kuo S, Johnston WM. Color accuracy of commercial digital cameras for use in dentistry. *Dent mater* 2006;22:553-559

15. Spear F. *The Art of Intra-Oral Photography*. Seattle Institute for Advanced Dental Education. 1999. p. 29-32.

16. Paul S, Peter A, Pietrobon N, Hammerle CH. Visual and spectrophotometric shade analysis of human teeth. *J Dent Res* 2002;81:578-92.

17. Ishikawa-Nagai S, Sato R, Furukawa K, Ishibashi K. Using a computer color matching system in color reproduction of porcelain restorations. part 1:Application of CCM to the opaque layer. *Int J Prosthodont* 1992;5:495-502.

18. Ragain JC. *Matching the Optical Properties of Direct Esthetic Dental Rstorative materials to Those of Human Enamel and Dentin*. A dissertation for the partial fulfillment of the requirement for the nf degree Doctor of Philosophy in the Graduate School of The Ohio State University: Columbus, OH; 1998.

19. Browning WD, Chan DC, Blalock JS, Brackett MS. A Comparison of Human Raters and an Intra-oral Spectrophotometer. *Oper Dent* 2009;34:337-343

20. Christopher CK Ho, BDS Hons (Syd) Shade Selection Aust. *Dental Prac*, sept/ oct 2007.

21. Da Silva JD, Park SE, Weber HP, Ishikawa-Nagai S. Clinical performance of a newly developed spectrophotometric system on tooth color reproduction. *J Prosth Dent*. 2008;99(5):361-368.

22. Yuan JC, Brewer JD, Monaco EA, Davis EL. Defining a natural tooth color space based on a 3-dimensional shade system. *J Prosthet Dent*. 2007;98(2):110–119.

23. Lagouvardos PE, Fougia AG, Diamantopoulou SA, PolyzoisGI. Repeatability and interdevice reliability of two portable color selection devices in matching and measuring tooth color. *J of Prosthet Dent*. 2009;101(1):40-45.

24. Kim-Pusateri S, Brewer JD, Davis EL, Wee AG. Reliability and accuracy of four dental shade-matching devices. *J Prosthet Dent*. 2009;101(3):193-199.