

**Evaluation of Immediate Loading of Dental Implants: A Review Article**

<sup>1</sup>Dr Yamini unni PT, Second year PG student, Department of Periodontics, Amrita School of Dentistry, Amrita Institute of Medical Sciences, Ponekkara, Kochi, Kerala, Pin-682041.

<sup>2</sup>Dr Karthik Prabhu N, First year PG student, Department of Periodontics, Amrita School of Dentistry, Amrita Institute of Medical Sciences, Ponekkara, Kochi, Kerala, Pin-682041.

<sup>3</sup>Dr Biju Balakrishnan, Reader, Department of Periodontics, Amrita School Of Dentistry, Amrita Institute of Medical Sciences, Ponekkara, Kochi, Kerala, Pin-682041.

<sup>4</sup>Dr Angel Fenol, Professor and HOD, Department of Periodontics, Amrita School of Dentistry, Amrita Institute of Medical Sciences, Ponekkara, Kochi, Kerala, Pin-682041.

<sup>5</sup>Dr Mohammed Shereef, Professor, Department of Periodontics, Amrita School of Dentistry, Amrita Institute of Medical Sciences, Ponekkara, Kochi, Kerala, Pin-682041.

<sup>6</sup>Dr Anuradha Bhaskar, Professor, Department of Periodontics, Amrita School of Dentistry, Amrita Institute of Medical Sciences, Ponekkara, Kochi, Kerala, Pin-682041.

**Corresponding author:** Dr Yamini unni PT, Second year PG student, Department of Periodontics, Amrita School of Dentistry, Amrita Institute of Medical Sciences, Ponekkara, Kochi, Kerala, Pin-682041.

**Citation of this Article:** Dr Yamini unni PT, Dr Karthik Prabhu N, Dr Biju Balakrishnan, Dr Angel Fenol, Dr Mohammed Shereef, Dr Anuradha Bhaskar, “Evaluation of Immediate Loading of Dental Implants: A Review Article”, IJDSIR- May - 2020, Vol. – 3, Issue -3, P. No. 15 – 21.

**Copyright:** © 2020, Dr.Mohd Athar Iqbal, et al. This is an open access journal and article distributed under the terms of the creative commons attribution noncommercial License. Which allows others to remix, tweak, and build upon the work non commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**Type of Publication:** Review Article

**Conflicts of Interest:** Nil

**Abstract**

The loss of tooth in the tasteful territory is frequently a horrible encounter for the patient. Patients may endure genuine or saw unfavorable impacts following the loss of at least one teeth. Dental implants offers the most financially savvy and long haul answer for supplanting of missing teeth with high normal future, furnishing the patient with the best feeling that all is well with the world and prosperity. As of late, immediate implant position after extraction of tooth with early loading has gotten

increasingly normal. The upsides of this system incorporate less careful mediations, decrease in general treatment time, diminished delicate and hard tissue misfortune, and mental fulfillment to the patient. immediately loaded implants help in balancing mental issues as the patient don't need to stay edentulous after position of implants as on account of traditionally loaded implants. Furthermore, they give capacity, solace, and discourse, therefore, immediately improved patient satisfaction.

## **Introduction**

Oral wellbeing and oral human services are imperative to keep up legitimate rumination, processing, phonation, appearance, and mental prosperity<sup>1</sup>. The loss of at least one teeth because of any explanation may antagonistically influence the oral wellbeing with an influenced appearance being the most genuine ramification for the patient and prime explanation referred to by the patient for looking for prosthodontic treatment<sup>2</sup>. Tooth misfortune can happen for an assortment of reasons which incorporates inborn non appearance, injury, illnesses of the dentition (e.g., caries or periodontal malady), just as mechanical disappointment. Tooth misfortune can likewise happen auxiliary or associatively to different fundamental maladies, for example, malignant growth, cardiovascular infection, diabetes mellitus, and osteoporosis. In this way, it is imperative to keep up great oral cleanliness, yet additionally generally speaking health<sup>3</sup>

Patients may endure genuine or saw negative impacts following the loss of at least one teeth. Mental impacts extend from insignificant to neuroticism. Tooth misfortune adds to loss of certainty, shirking of giggling out in the open, hesitance to frame cozy relationship, particularly when front teeth are missing<sup>4</sup> Most developed approach to supplant missing teeth is dental embed which is intended to imitate the characteristic tooth root and crown of the normal tooth. This technique safeguards the gingival mucosa and bone with no harm to adjoining teeth. Ordinary methodology for embed situation includes extraction of culpable tooth, trusting that extraction attachment will mend, inclusion of embed, and again sitting tight for 3–6 months for reconciliation of embed with encompassing bone; after this system, another medical procedure is important to uncover the embed and to put a prosthetic abutment.<sup>5</sup> Taking into thought the

prosthetic treatment, the patient needed to hold up to 8 a year for a lost tooth to be supplanted. Due to these deficiencies identified with ordinary method, systems were created to generously abbreviate the whole treatment by situation of implant following extraction of tooth followed by quick loading of implant with prosthesis<sup>6</sup> several long haul contemplates showed high achievement rates when an exacting careful and prosthodontic convention was followed<sup>6,7</sup>. One of the key criteria for appropriate osseointegration was the requirement for emptied submerged recuperating of the implants for a time of three to a half year (two-organize approach)<sup>6,7</sup>. The worry was that untimely loading would cause micromotion prompting sinewy exemplification of the implant. In any case, the long haul achievement and consistency with dental inserts urged clinicians to reevaluate the first Brånemark convention, since it depended fundamentally on clinical perceptions and not really on biologic principles<sup>8</sup>. The delayed postoperative period following implant position prompted unavoidable troubles in persistent administration. During this osseointegration period, a few inconveniences, including free false teeth, cracked prosthesis, sore spots and intermittent temporary relines, tormented the clinician and the patient. What's more, the need for a removable prosthesis, in any event, for a brief period, was an obstruction in itself for certain patients whose essential objective was to stay away from a removable dental replacement in any case. Along these lines, the idea of submerged recuperating was tested first by Schroeder and afterward in creature and human examinations by Ericsson et al, also, Becker et al. who exhibited that one-and two-arrange approaches both prompted likewise victories.

### What Is Immediate Loading...?

The logical writing is overflowing with meanings of immediate loading of dental implants. Misch et al., in 2004,<sup>13</sup> offered a few arrangements of immediate loading:

- Immediate occlusal loading alludes to full useful occlusal loading of an implant inside about fourteen days of arrangement.
- Early occlusal loading alludes to practical loading between about fourteen days and 3 months of implant arrangement.
- Nonfunctional prompt reclamation alludes to implant prostheses set inside about fourteen days of embed situation with no direct utilitarian occlusal loading.
- Nonfunctional early reclamation alludes to embed prostheses conveyed between about fourteen days and 3 months from embed arrangement.
- Delayed occlusal loading alludes to the reclamation of an embed over 3 months after arrangement.

These classes help to portray the time span of the therapeutic period of embed medical procedure. In 2006, Wang et al.<sup>14</sup> gave a definition dependent on an accord from the International Congress of Oral Implantologists (Upper Montclair, NJ) in which quick stacking was depicted as a system in which the embed bolstered reclamation is set into utilitarian occlusal loading inside 48 hours of embed inclusion. Besides, a differentiation was made between the prompt reclamation for tasteful purposes, in which the rebuilding was set out of occlusal contacts, and genuine quick loading.

### Principles of Immediate Loading

It has been portrayed by Frost mechanostat hypothesis which recommends that bone adjusts by various biologic procedures:<sup>15</sup> trivial, physiological, over-burden and obsessive. Renovating is portrayed as a synchronous procedure of arrangement and resorption that replaces already existing bone, tends to remove or save bone and is

enacted by decreased mechanical utilization in the minor loading zone or miniaturized scale harm in the neurotic loading zone. With the assistance of the implant at whatever point a controlled burden is applied deep down, bone reacts by renovating its design as indicated by the greatness and the heading of the heap applied.

Fundamental goal of immediately loaded implant prosthesis is to lessen the danger of occlusal over-burden and accordingly, bringing about increment in the redesigning pace of bone. Woven and lamellar are the two sorts of bone shaping at the interface. Woven bone is created in light of phenomenal loading condition, framing at a pace of in excess of 60 microns every day and is seen as less mineralized<sup>12</sup> while lamellar bone structures at a pace of 1-5 microns every day. Subsequently, a higher turnover rates lead to higher dangers for the bone-embed interface.<sup>16</sup>

### Implant Loading Time Protocols

Esposito et al.<sup>17</sup> have characterized 3 conventions for implant load timing: immediate loading implants (ILI), inside multi week from embed arrangement; early loading implants (ELI), between multi week and 2 months; and conventional loading implants (CLI), following 2 months from embed situation.

Two sub classifications bring up the distinctive stacking methodology:

- 1) Occlusal loading or Non Occlusal loading
- 2) Direct loading or Progressive loading.

This Cochrane methodical survey<sup>18</sup> presumes that there is no persuading proof regarding a clinically significant contrast in prosthesis disappointment, embed disappointment, or bone misfortune related with various stacking times of inserts.

Additionally ILI, ELI and CLI were found not fundamentally unique as far as the related negligible bone misfortune, changes in embed strength, and wellbeing

status of the peri-embed tissues, which showed that these stacking conventions acted likewise once osseointegration occurred<sup>19</sup>.

### Signs and Contraindications

Only one out of every odd patient or each tooth site is demonstrated for the immediate loading approach. Patients must comprehend the impediments of such treatment and be happy to acknowledge the deductively based careful steps.<sup>20</sup> Chief among them is the way that, so as to restrict the useful powers during osseointegration, patients need to refrain from biting anything other than delicate nourishment or in any case applying power to the reclamation for around 3 months.<sup>21</sup>

### Indications

- Completely edentulous jaw.
- Partially edentulous jaw.
- Patients with missing dentition requiring long range fixed fractional dental replacement.
- Patient who are not ready to utilize a removable sort prosthesis<sup>20,23,25</sup>.
- Immediate loading convention ought to be restricted to the patients who have the most to pick up and least to lose. Eg. Patients who can't endure a removable prosthesis because of social or mental reasons.
- Patients who can't hang tight for 3 months for the prosthesis.

Different signs depicted are as follows<sup>23</sup>

- Poor oral solid coordination.
- Ridiculous patient desires for complete dentures.
- Tolerant mentally against removable prosthesis<sup>4</sup>.
- Single tooth misfortune; keep away from planning of sound teeth.

### Contraindications

The proposed contraindications, in general, for consideration of a prompt occlusal convention incorporate the accompanying<sup>20,21</sup>:

- Chronic smoker.
- If bone volume isn't sufficient.
- If density of bone isn't acceptable (D4).
- Parafunctional; biting propensities (bruxing, grasping, tongue push).

### Primary Stability

Micromovements have been considered, since the beginning of embed dentistry, one of the primary hazard for the accomplishment of osseointegration<sup>24</sup>. It has been demonstrated that if the micromovements extend results to be more than 150  $\mu\text{m}$ , this could imperil the osseointegration procedure. This extreme micromotion results to be straightforwardly ensnared in the development of the implant stringy exemplification<sup>25,26</sup>. The writing proposes that there is a basic edge of micromotion above which sinewy exemplification beats osseointegration. This basic level, in any case, was not zero micromotion as for the most part deciphered. Rather, the endured micromotion limit was found to lie somewhere close to 50 and 150 microns<sup>24, 27</sup>. Right now run an early burden on the embed surface could even animate the recently framed issue that remains to be worked out, quickening the osseointegration procedure. Essential implant strength is impacted by numerous components including nearby bone quality and amount, implant large scale plan and careful procedure.

### Complications Associated and Solutions

The ILI convention all the more frequently prompts specialized entanglements. The most widely recognized of those were cracks of the prostheses, releasing of the projection screws and dental replacement shaping modifications. The last could be clarified by the auxiliary

gingival recuperating after medical procedure in the early loading prostheses, which may bring about space around the projections, while relining impressions acted in the regular loading inserts after a time of mending maintained a strategic distance from this space<sup>28</sup>.

Clinical entanglements which are related with immediate implants are slackening and crack of screw, breaks of prosthesis and veneering materials, proceeding with minimal bone misfortune underneath the principal string along the implant, implant cracks, and implant misfortune. To conquer such complexities, following techniques dependent on utilization of sound biomechanical standards, for example, uninvolved attack of the prosthesis, cantilever length can be decreased, narrowing the bucco-lingual/mesiodistal interface and embed prosthesis, to keep up embed load inside the physiological furthest reaches of individualized impediment, lastly to give long haul strength of implants and implant prostheses.<sup>29</sup> IL of dental inserts has as of late picked up prevalence and have yielded a wide scope of clinical survival.<sup>30-34</sup> It has been reported that utilizing these inserts direct bone interface, every so often, could be created and kept up for more than 20 years.<sup>40</sup> In the center 1980s, high achievement rates from immediately loaded implants had been first recorded in quite a while exactly when the arrange embed convention became popular. Babbush et al. (1986) revealed a combined achievement pace of 88% on 1739 immediate loading TPS implants.<sup>41</sup> Furthermore, it is suggested that uncommon careful methods be utilized to expand bone thickness in the embedded before implant inclusion to improve essential dependability which is considered as best quality level to diminish the micromovement and well as to set up long haul accomplishment for prompt loading.<sup>34</sup> It is deductively demonstrated that IL is an attractive methodology, if the result regarding implant endurance

and achievement is similar with that of regular loading yet at the same time Future investigations ought to be done to assess long haul information of immediate rebuilding efforts on implants and potential utilizations of this system in circumstances where nature of bone is poor, different inserts or expansion strategies exist, must be overcome.

### **Conclusion**

The most recent decade has seen a significant move in implant dentistry from the protracted recuperating unique conventions to immediate loading has requested a psychological upheaval. Immediate loading of a dental implant comprises of a nonsubmerged organize medical procedure in which loading of the implant is finished with a temporary rebuilding at a similar arrangement or presently. immediate loading of dental implants are demonstrated to diminish the treatment time and in this way increment persistent acknowledgment. Future investigations ought to be directed to assess long haul information of quick reclamations on implants and potential uses of this method in circumstances where issues of poor bone quality, different implants or expansion systems must be survived.

The arrangement of implants and their quick rebuilding, regardless of whether temporary or final, can be invaluable. Nonetheless, mind and fitting careful and prosthetic contemplations should be profoundly mulled over when playing out these strategies. As we play out these systems, we're expanding the likelihood for intricacies as more parts of medicines are being rendered. Its prosperity rate may contrast somewhat from finishing strategies in a progressively ordinary manner. Of the best and most invaluable utilization of prompt reclamation of inserts are those cases wherein tasteful needs and delicate tissue protection are generally significant.

## Reference

1. Meyer MS, Joshipura K, Giovannucci E, Michaud DS. A review of the relationship between tooth loss, periodontal disease, and cancer. *Cancer Causes Control* 2008;19:895-907.
2. Desvarieux M, Demmer RT, Rundek T, Boden-Albala B, Jacobs DR Jr., Papapanou PN, et al. Relationship between periodontal disease, tooth loss, and carotid artery plaque: The oral infections and vascular disease epidemiology study (INVEST). *Stroke* 2003;34:2120-5.
3. Anil S, Preethanath RS, AlMoharib HS, Kamath KP, Anand PS. Impact of osteoporosis and its treatment on oral health. *Am J Med Sci* 2013;346:396-401.
4. Craddock HL. Consequences of tooth loss: 1. The patient perspective – Aesthetic and functional implications. *Dent Update* 2009;36:616-9.
5. Schropp L, Isidor F. Timing of implant placement relative to tooth extraction. *J Oral Rehabil* 2008;35 Suppl 1:33-43.
6. Singh A, Gupta A, Yadav A, Chaturvedi TP, Bhatnagar A, Singh BP. Immediate placement of implant in fresh extraction socket with early loading. *Contemp Clin Dent* 2012;3 Suppl 2:S219-22.
6. Brånemark PI, Hansson BO, et al, Osseointegrated implants in the treatment of the edentulous jaw. Experience from a 10-year period. *Scand J Plast Reconstr Surg (Suppl)* 16:1-132, 1977.
7. Adell R, Lekholm U, et al, A 15-year study of osseointegrated implants in the treatment of the edentulous jaw. *Int J Oral Surg* 10:387-416, 1981.
8. Albrektsson T, Direct bone anchorage of dental implants. *J Prosth Dent* 50:255-61, 1983.
9. Szmukler-Moncler S, Piattelli A, Favero GA, et al, Considerations preliminary to the application of early and immediate loading protocols in dental implantology. *Clin Oral Implants Res* 11:12-25, 2000.
10. Schroeder A, Van Der Zypen E, Stich H, et al, The reaction of bone, connective tissue, and epithelium to endosteal implants with titanium-sprayed surfaces. *J Oral Maxillofac Surg* 9:15-25, 1981.
11. Ericsson I, Nilner K, Klinge B, Glantz PO, Radiographic and histological characteristics of submerged and non-submerged titanium implants. An experimental study in the Labrador dog. *Clin Oral Implants Res* 7:20-6, 1996.
12. Ericsson I, Randow K, et al, Some clinical and radiographic features of submerged and non submerged titanium implants. A five-year followup study. *Clin Oral Implants Res* 8(5):422-6, 1997.
13. Misch CE, Wang HL, Misch CM, et al : Rationale for the application of immediate load in implant dentistry: part 1, *Implant Dent* 13 : 207 - 217 , 2004 .
14. Wang HL, Ormianer Z, Palti A, et al : Consensus conference on immediate loading: the single tooth and partial edentulous areas, *Implant Dent* 15 : 324 - 333 , 2006
15. Mahesh P, Chalapathi Kumar VH, Saran Babu KA. Immediate loading of dental implants: a review-Part I. *Annals and Essences of Dentistry* 2014;6(2):44-50.
16. Romanas GE et al. Histologic and Histomorphometric Evaluation of Peri-implant Bone Subjected to Immediate Loading: An Experimental Study with *Macaca Fascicularis*. *Int J Oral Maxillofac Implants* 2002;17:44-51
17. Esposito M, Grusovin MG, Maghairy H, Worthington HV. Interventions for replacing missing teeth: different times for loading dental implants. *Cochrane Database Syst Rev*. 2013;3:CD003878.
18. Enriquez-Sacristan C, Barona-Dorado C, Calvo-Guirado JL, Leco-Berrocal I, Martinez-

- Gonzalez JM. Immediate post-extraction implants subject to immediate loading: a meta-analytic study. *Med Oral Patol Oral Cir Bucal*. 2011;16:e919-24.
19. Sanz-Sanchez I, Sanz-Martin I, Figuro E, Sanz M. Clinical efficacy of immediate implant loading protocols compared to conventional loading depending on the type of the restoration: a systematic review. *Clin Oral Implants Res*. 2015;26:964-82. 52.
- Zhu Y, Zheng X, Zeng G, Xu Y, Qu X, Zhu M, Lu E. Clinical efficacy of early loading versus conventional loading of dental implants. *Sci Rep*. 2015; 5:15995.
20. Mahesh P,Chalapathi Kumar VH , Saran Babu KA. Immediate loading of dental implants: a review-Part I. *Annals and Essences of Dentistry* 2014;6(2):44-50
21. Evans GH, Mendez J, Caudil RF. Loaded and Non loaded Titanium versus Hydroxyapatite- Coated Threaded Implants in the Canine Mandible. *Int J Oral Maxillofac Implants*.1996; 11:360–371.
22. Misch CE: Workshop Guidelines On Immediate Loading In Implant Dentistry. *J Oral Imp* 2004;30(5):283-8.
23. Hobo S, Ichida E, Gracia LT. *Osseointegration And Occlusal Rehabilitation*: Quintessence Publishing Co., 1989
24. Szmukler-Moncler S, Piattelli A, Favero GA, Dubruille JH. Considerations preliminary to the application of early and immediate loading protocols in dental implantology. *Clin Oral Implants Res*. 2000;11:12-25.
25. Szmukler-Moncler S, Salama H, Reingewirtz Y, Dubruille JH. Timing of loading and effect of micromotion on bone-dental implant interface: review of experimental literature. *J Biomed Mater Res*. 1998;43: 192-203.
26. Papaspyridakos P, Chen CJ, Chuang SK, Weber HP. Implant loading protocols for edentulous patients with fixed prostheses: a systematic review and meta-analysis. *Int J Oral Maxillofac Implants*. 2014; 29 Suppl:256-70.
27. Javed F, Ahmed HB, Crespi R, Romanos GE. Role of primary stability for successful osseointegration of dental implants: Factors of influence and evaluation. *Interv Med Appl Sci*. 2013;5:162-7.
28. JayeshRaghvendra S., Dhinakarsamy. *J Pharm Bioallied Sci*. 2015 Apr; 7 (Suppl1): S226–S229.
29. Strock AE, Strock M. Experimental work on a method for the replacement of missing teeth by direct implantation of a metal support into the alveolus. *Am J Orthod Oral Surg*. 1939;25:467–472.
30. Linkow LI. The blade-vent—a new dimension in endosseous implants. *Dent Concepts*. 1968;11:13–18.
31. Cranin AN, Rabkin MF, Garfinkel L. A statistical evaluation of 952 endosteal implants in humans. *J Am Dent Assoc*. 1977;94:315–320.
32. Smithloff M, Fritz ME. The use of blade implants in a selected population of partially edentulous adults: a fiveyear report. *J Periodontol*. 1976;47:19–24.
33. Kapur KK. Veterans Administration co-operative dental implant study comparison between fixed partial dentures supported by Blade-Vent implants and partial dentures.*J Prosthet Dent*. 1987;59:499–512.
34. Linkow LI, Donath K, Lemons JE. Retrieval analysis of a blade implant after 231 months of clinical function.*Implant Dent*. 1992;1:37–43.
35. Babbush, C. A., Kent, J. N., Misiek, D. J. Titanium plasma-sprayed (TPS) screw implants for the reconstruction of the edentulousmandible. *Journal of Oral & Maxillofacial Surgery* 1986, 44: 274–282.
36. Romanos GE. Present Status of Immediate Loading of Oral Implants.*J Oral Implantol* 2004;30:189-97.